

General

Title

Bipolar disorder: the percentage of patients with Bipolar I Disorder with mania/hypomania, mixed or cycling symptoms and behaviors who have evidence of use of pharmacotherapy agent with antimanic properties during the first 12 weeks of treatment.

Source(s)

STABLE (STAndards for BipoLar Excellence) performance measures. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. various p.

Measure Domain

Primary Measure Domain

Process

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the [Measure Validity](#) page.

Secondary Measure Domain

Does not apply to this measure

Brief Abstract

Description

This measure is used to assess the percentage of patients with Bipolar I Disorder with mania/hypomania, mixed or cycling symptoms and behaviors who have evidence of use of a pharmacotherapy agent with antimanic properties during the first 12 weeks of treatment.

Rationale

Goals of Acute Treatment: Mania or Mixed Episodes

If present, a rapid response to agitation, aggression and impulsivity is desired.

Remission of symptoms with a return to usual levels of psychosocial functioning.

Initial Treatment Considerations

Treatment selection is dependent on illness severity, associated clinical features such as rapid cycling or psychosis, and patient preference.

Treatment choice should also consider patient history, potential side effects and individual therapeutic response to specific pharmacotherapy agents.

Initial Treatment Recommendations

Guidelines suggest lithium or valproate (divalproex/divalproex sodium/valproate sodium/valproic acid) alone or in combination with an antipsychotic; alternatives to lithium or valproate may include carbamazepine.

Clozapine has been noted as an effective antimanic therapy in treatment-resistant bipolar disorder; however, it is recommended with caution due to safety concerns and requires monitoring.

Primary Clinical Component

Bipolar I Disorder; depression; antimanic agent

Denominator Description

Patients with Bipolar I Disorder episodes with

Manic/hypomanic symptoms or behaviors

Mixed symptoms or behaviors

Cycling symptoms or behaviors

See the related "Denominator Inclusions/Exclusions" field.

Numerator Description

Patients with evidence of use of an antimanic agent during the first 12 weeks of pharmacotherapy treatment (see the related "Numerator Inclusions/Exclusions" field)

Evidence Supporting the Measure

Evidence Supporting the Criterion of Quality

A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence

A formal consensus procedure involving experts in relevant clinical, methodological, and organizational sciences

One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

Evidence Supporting Need for the Measure

Need for the Measure

Unspecified

State of Use of the Measure

State of Use

Current routine use

Current Use

Internal quality improvement

Application of Measure in its Current Use

Care Setting

Ambulatory Care

Behavioral Health Care

Physician Group Practices/Clinics

Professionals Responsible for Health Care

Advanced Practice Nurses

Physicians

Psychologists/Non-physician Behavioral Health Clinicians

Lowest Level of Health Care Delivery Addressed

Individual Clinicians

Target Population Age

Age greater than or equal to 18 years

Target Population Gender

Either male or female

Stratification by Vulnerable Populations

Unspecified

Characteristics of the Primary Clinical Component

Incidence/Prevalence

Unspecified

Association with Vulnerable Populations

Unspecified

Burden of Illness

Unspecified

Utilization

Unspecified

Costs

Unspecified

Institute of Medicine (IOM) Healthcare Quality Report Categories

IOM Care Need

Getting Better

Living with Illness

IOM Domain

Effectiveness

Data Collection for the Measure

Case Finding

Users of care only

Description of Case Finding

Patients with diagnosis of Bipolar I Disorder; manic/hypomanic, mixed or cycling: International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) or Diagnostic and Statistical Manual of Mental Disorders, Fourth Revision, Text Revision (DSM-IV-TR): 296.0x; 296.1x; 296.4x; 296.6x

Denominator Sampling Frame

Patients associated with provider

Denominator Inclusions/Exclusions

Inclusions

Patients 18 years of age or older with an initial diagnosis or new presentation/episode of bipolar disorder

AND

Documentation of Bipolar I Disorder with manic/hypomanic; mixed; or cycling symptoms or behaviors; to include at least one of the following:

Codes 296.0x; 296.1x; 296.4x; 296.6x documented in body of chart, such as a pre-printed form completed by a clinician and/or codes documented in chart notes/forms

Diagnosis or impression documented in chart indicating Bipolar I Disorder with current symptoms or behaviors of mania/hypomania; mixed state symptoms or behaviors; or indications of cycling

Exclusions

Exclude case from denominator population if numerator-required pharmacotherapy is not prescribed and the reason(s) for not prescribing is documented in chart, such as, not clinically indicated; not prescribed for related medical reasons (allergy, drug interaction); or not prescribed for patient reasons (patient refusal), etc.

Relationship of Denominator to Numerator

All cases in the denominator are equally eligible to appear in the numerator

Denominator (Index) Event

Clinical Condition

Encounter

Denominator Time Window

Time window is a single point in time

Numerator Inclusions/Exclusions

Inclusions

Documented treatment with at least one pharmacotherapy agent with antimanic properties (Refer to the "Data Dictionary Reference" in the original measure documentation for specified medications.)

AND

Timeframe:

Documentation of the bipolar disorder pharmacotherapy must be within the first 12 weeks of treatment.

Exclusions

Unspecified

Measure Results Under Control of Health Care Professionals, Organizations and/or Policymakers

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

Numerator Time Window

Fixed time period

Data Source

Administrative data

Medical record

Level of Determination of Quality

Individual Case

Pre-existing Instrument Used

Unspecified

Computation of the Measure

Scoring

Rate

Interpretation of Score

Better quality is associated with a higher score

Allowance for Patient Factors

Unspecified

Standard of Comparison

Internal time comparison

Evaluation of Measure Properties

Extent of Measure Testing

The STABLE measures were developed using the RAND Appropriateness Method and have been shown to have content validity and face validity.

Data feasibility testing was performed to determine the availability of the data elements required in the measure numerator and denominator specifications.

Inter-abtractor reliability testing was performed to assess the data collection strategy. The data collection strategy included data collection forms; data dictionary references and abtractor instructions.

A field study was conducted to determine measure conformance in an appropriate convenience sample.

Refer to the references listed below for further information.

Evidence for Reliability/Validity Testing

STABLE performance measures: data feasibility testing & results. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. 2 p.

STABLE performance measures: development process & validity ratings. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. 3 p.

STABLE performance measures: field study process & conformance findings. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. 3 p.

STABLE performance measures: inter-abtractor reliability testing & results. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. 2 p.

Identifying Information

Original Title

Bipolar disorder: use of antimanic agent in BD I with mania/hypomania, mixed, or cycling.

Measure Collection Name

Standards for Bipolar Excellence (STABLE) Performance Measures

Submitter

Center for Quality Assessment and Improvement in Mental Health - Clinical Specialty Collaboration

Developer

STABLE Project National Coordinating Council - Clinical Specialty Collaboration

Funding Source(s)

AstraZeneca LLP, Wilmington, Delaware, provided financial sponsorship for the STABLE Project. They did not otherwise participate in the development of either the measures or toolkit.

Composition of the Group that Developed the Measure

The STABLE [National Coordinating Council \(NCC\)](#) was comprised of national

experts in bipolar disorder, psychiatry, primary care, and performance improvement. The NCC guided and directed the STABLE Project. NCC members agreed to serve with the understanding that the STABLE Performance Measures and Resource Toolkit would be fully transparent and available without cost in the public domain.

EPI-Q, Inc. , is a consulting company providing practice-based outcomes research, pharmacoeconomic studies, and quality improvement services. EPI-Q managed the STABLE Project.

Financial Disclosures/Other Potential Conflicts of Interest

Unspecified

Adaptation

Measure was not adapted from another source.

Release Date

2007 Jan

Measure Status

This is the current release of the measure.

The STABLE Project National Coordinating Council reaffirmed the currency of this measure in November 2010.

Source(s)

STABLE (STAndards for BipoLar Excellence) performance measures. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007. various p.

Measure Availability

The individual measure, "Bipolar Disorder: Use of Antimanic Agent in BD I with Mania/Hypomania, Mixed, or Cycling," is published in "STABLE (STAndards for BipoLar Excellence) Performance Measures." This document is available in Portable Document Format (PDF) from the [Center for Quality Assessment and Improvement in Mental Health \(CQAIMH\) Web site](#) .

Companion Documents

The following is available:

STABLE National Coordinating Council Resource Toolkit Workgroup. STABLE resource toolkit. Boston (MA): Center for Quality Assessment and Improvement in Mental Health; 2007 Mar. 67 p. This document is available in Portable Document Format (PDF) from the [Center for Quality Assessment and Improvement in Mental Health \(CQAIMH\) Web site](#) .

NQMC Status

This NQMC summary was completed by ECRI Institute on January 10, 2008. The information was verified by the measure developer on April 14, 2008. The information was reaffirmed by the measure developer on November 1, 2010.

Copyright Statement

The STABLE materials published on this Web site are copyrighted by CQAIMH. They may be used for research, teaching, and quality measurement/improvement activities " provided the following:

The materials are not sold, distributed or licensed for commercial purposes.
CQAIMH's copyright is acknowledged in reproductions of these materials.
Modifications to the materials are not made without CQAIMH's permission.

Disclaimer

NQMC Disclaimer

The National Quality Measures Clearinghouse" (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the [NQMC Inclusion Criteria](#).

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. Moreover, the views and opinions of developers or authors of measures represented on this site do not necessarily state or reflect those of NQMC, AHRQ, or its contractor, ECRI Institute, and inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.